

Resolving size distribution of black carbon internally mixed with snow: impact on snow optical properties and albedo

Cenlin He^{a,b,*}, Kuo-Nan Liou^b, and Yoshi Takano^b

^a*Advanced Study Program, National Center for Atmospheric Research, Boulder, CO 80301, USA*

^b*Joint Institute for Regional Earth System Science and Engineering and Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, CA 90095, USA*

**Presenting author (cenlinhe@ucar.edu)*

We develop a stochastic aerosol-snow albedo model that explicitly resolves the size distribution of aerosols internally mixed with various snow grains, based on a geometric-optics surface-wave (GOS) approach. We use the model to quantify effects of black carbon (BC) size distribution on snow albedo and optical properties for BC-snow internal mixing. Results show that BC-induced snow single-scattering coefficient enhancement and albedo reduction decrease by a factor of 2–3 with increasing BC effective radii from 0.05 to 0.25 μm , while polydisperse BC results in up to 40% smaller visible single-scattering coefficient enhancement and albedo reduction, compared to monodisperse BC with equivalent effective radii. We further develop parameterizations for BC size effects for application to climate models. Compared with a realistic polydisperse assumption and observed shifts to larger BC sizes in snow, respectively, assuming monodisperse BC and typical atmospheric BC effective radii inside snow grains could lead to overestimates of ~24% and ~40% in BC-snow albedo forcing averaged over different BC and snow conditions.

Preferred mode of presentation: Oral